



PREVENTING MALARIA ATOVACUONA-PROGUANIL

To avoid malaria in the area you visit, take ATOVACUONE-PROGUANIL.

Recommended dosage: 1 tablet **daily** (for people over 40 kg body weight) taken with food, preferably fatty, and plenty of liquid, or whole milk, after breakfast. Allow one hour before going to bed or lying down:

- Start 24 to 48 hours **before** entering a malaria-endemic geographic area.
- Continue **daily** during your **stay**.
- Continue **daily** for seven days **after** you leave the area. If you experience vomiting during the first hour after taking it, take another.

Side effects: The most frequent are headaches and an upset stomach.

Contraindications: Hypersensitivity to the active ingredient or excipients in the medicine or severe renal insufficiency. Do not take it together with Rifampicin, tetracyclines or Metoclopramide. There are no studies on its use during pregnancy. Avoid taking it during lactation.

REMEMBER:

No antimalarial prophylaxis ensures complete protection, so the first defence against the disease is to avoid mosquito bites.

Pregnant women and children are especially susceptible to severe malaria.

If you experience fever, joint pain, general malaise or diarrhoea from one week after entering the risk area or up to three months after leaving it, consult a doctor to rule out the possibility of malaria.

If you experience severe side effects, discontinue the medication and seek medical advice.

Read the medicine's patient package insert carefully.